## Kertas Asli/Original Articles

Parent-Implemented Language Intervention for Late Talkers: A Scoping Review (Intervensi Bahasa Dilaksanakan Ibu Bapa untuk Kanak-kanak Lambat Bercakap: Satu Tinjauan Skop)

#### ABSTRACT

Parent-implemented language intervention (PILI) is one of the intervention approaches used in managing late talkers (LTs). Yet, there are few evidence-based PILI programs available for parents of LTs. This scoping review aimed to (a) assess the characteristics of participants depicted in the literature related to PILI programs for parents of LTs, (b) provide an overview of the structure and design of the available PILI programs for parents of LTs, and (c) explore the effectiveness of the programs in relation to the characteristics of reviewed studies. The scoping review was performed by adhering to the general principles prescribed by Arksey and O'Malley (2005). Articles that fulfilled the inclusion criteria and were published from 1980 to 2018 were selected. Two reviewers independently charted the information from the identified articles. A total of 15 articles were selected. The results were reviewed in terms of participant characteristics, intervention characteristics, and effectiveness of PILI programs. Most adult participants in PILI studies were mothers and included child participants with either expressive language delay or receptive and expressive language delay. Moreover, there were different structures and designs of PILI programs for parents of LTs. Preliminary evidence indicated that PILI was more effective than no/delayed intervention and could be as effective as direct therapy provided by clinicians. A discussion related to the findings was also presented.

Keywords: parent-implemented language intervention, late talkers, parent training, language intervention

### ABSTRAK

Intervensi bahasa yang dilaksanakan oleh ibu bapa (PILI) merupakan salah satu pendekatan intervensi yang digunakan ketika menguruskan kes kanak-kanak lambat bercakap (LTs). Terdapat beberapa program PILI yang boleh digunakan untuk melatih ibu bapa yang mempunyai anak LTs. Tujuan tinjauan skop ini dijalankan adalah untuk; a) menilai ciriciri peserta yang menyertai kajian berasaskan PILI, b) memberi gambaran tentang struktur dan reka bentuk program-program PILI yang ada khusus untuk ibu bapa yang mempunyai anak LTs dan c) meneroka keberkesanan program-program PILI. Tinjuan skop ini dijalankan berdasarkan garis panduan yang diperkenalkan oleh Arksey dan O'Malley (2005). Artikel-artikel yang memenuhi kriteria dan diterbitkan di antara tahun 1980 hingga 2018 dipilih. Pada peringkat akhir, 15 artikel telah dipilih. Dua orang penilai mencartakan maklumat-maklumat berkaitan daripada artikel-artikel yang telah dipilih secara berasingan. Maklumat-maklumat yang diekstrak daripada artikel-artikel ini adalah ciri-ciri peserta dan intervensi dan keberkesanan program-program PILI. Kebanyakan peserta dewasa di dalam kajian PILI adalah ibu dan peserta kanak-kanak mempunyai kelambatan bahasa ekspresif sahaja atau ekspresif dan kefahaman. Selain daripada itu, terdapat pelbagai reka bentuk dan struktur program PILI. Bukti awal menujukkan bahawa PILI adalah lebih berkesan daripada tiada/melewatkan intervensi dan efektif sama seperti terapi yang diberikan secara terus. Perbincangan berkaitan dengan penemuan tinjaun skop ini juga disertakan di dalam artikel ini.

Kata kunci: intervensi bahasa dilaksanakan ibu bapa, kanak-kanak lambat bercakap, latihan ibu bapa, intervensi bahasa

## INTRODUCTION

#### LATE TALKERS (LTS)

Late talkers (LTs) refer to toddlers who acquire language slower than their typically developing peers, despite normal cognition, sensory, motor, and neurological systems (Rescorla 2009). The age of LTs described in the literature varies from 18 to 42 months (Cable & Domsch 2011; Deveney et al. 2017; Hawa & Spanoudis 2014; Rescorla 2009; Roberts & Kaiser 2012). Desmarais and colleagues (2008) highlighted two sets of criteria available in the literature to describe LTs. Although there is a similarity in the cause of the problem, there is a fine line distinguishing these criteria sets. In the first classification, the criteria are more restrictive. To be identified as LT, a child needs to have limited expressive vocabulary, characterized by either vocabulary size falls below the 10th percentile of a normative sample (Henrichs et al. 2011) or less than 50 words or no word combination at the age of two, with intact comprehension abilities (Rescorla 1989). On the contrary, the second classification is much broader than the first, as it only includes the criterion of limited expressive vocabulary mentioned above but not the comprehension abilities part (e.g., Horwitz et al. 2003). Therefore, a larger proportion of children can meet the criteria, including children with comprehension problems. Other terms used to describe the conditions mentioned above are 'early language delay' (Scarborough & Dobrich 1990), 'late language emergence' (Zubrick et al. 2007), and 'expressive language delay' (Hawa & Spanoudis 2014).

For children whose language delay is secondary to developmental deficits, further assessment or intervention is integral. In the case of LTs, appropriate clinical management is still debated. Upon anticipating that most LTs outgrow the issue at hand, the 'watch and see' approach has been prescribed, where the LTs are monitored regularly (Paul 2001). The main reason for this recommendation is that although language problems are not resolved for some LTs by the time they reach school age, they are able to keep up with the curriculum demands and will not suffer serious academic difficulties. The reason is in accord with that reported by Dale et al. (2014). They discovered that recovered LTs possess comparable language and reading abilities at ages 7 and 12, although the mean for language measures for the recovered group was below the mean for the total sample. In addition, Rescorla (2009) revealed that at the age of 17, LTs and typical comparison children in her study did not differ significantly in mathematics, reading, and writing abilities.

Despite the abovementioned research findings that indicated that LTs might not suffer from significant

academic difficulties, LTs are at risk for language disorder. Leonard (2014) stated that at least one in five LTs would be diagnosed with developmental language disorder (DLD) as their language problems persisted until school age and could not be linked to any biomedical etiologies. Children with DLD usually have morphosyntax, semantic, and pragmatic problems, which are crucial for social and academic success (Kaderavek 2011). The finding of the longitudinal study by Armstrong et al. (2016), which investigated educational, employment, and mental health outcomes of young adults with and without a history of language problems, showed that at the age of 21 years old, adults with a history of deteriorated or persistent language problems were less likely to engage in education, employment, or training, including apprenticeships. They also exhibited a greater risk for alcohol and substance abuse/misuse, as well as affective disorders. In a similar vein, St Clair et al. (2019) emphasized that children at risk for DLD had increased levels of emotional difficulties at 11 years of age compared to the general population group. Clinically, the aforementioned arguments highlighted the significance of early identification and intervention of LTs.

## INTERVENTIONS FOR LTS

One important factor that may affect the outcome of intervention for children with primary language problems is suitable service delivery (Law et al. 2017). Ebbels et al. (2019) discussed three different tiers in speech-language therapy service delivery for children with language problems: Tier 1, Tier 2, Tier 3A, Tier 3B. Tier 1 may involve training other professionals and conducting parenting education programs to promote the development of speech, language, and communication in general. Tier 2, 3A, and 3B focus on intervention for children at risk or with language problems. Therefore, intervention for LTs can be spread across Tier 2, 3A, and 3B where parental group training (Tier 2), individual parental training (Tier 3A), and direct clinician-led intervention (Tier 3B) can be administered. However, changes in the clinical landscape can be seen as indirect approaches such as parental training are more preferred in younger children (Law et al. 2019). This transition can be linked with the general awareness that children's environment is associated with their language learning abilities and recommendations by the policy drivers (Deveney et al. 2017; Law et al. 2019; Roberts & Kaiser 2011). Due to that, this study will only focus on Tier 2 and Tier 3A, where parent-implemented language intervention (PILI) is applied.

The PILI is underpinned by principles of familycentred care (Espe-sherwindt & Serrano 2016). Based on this approach, therefore, the role of speech-language pathologists (SLPs) reflects more as parents' educators, facilitators, and/or consultants (American Speech-Language-Hearing Association [ASHA] 2008) to empower parents to be the primary interventionist via structured teaching programs and coaching (see Buschmann *et al.* 2009, Whitehurst *et al.* 1991). A recent study reported that 45% of 4020 SLPs conducted PILI when managing children with language problems using evidence-based or non-evidence-based programs (Law *et al.* 2019). They spent 12.70% of their time training parents besides training professionals, conducting assessments and direct therapy, and doing other administration works (Pring et al. 2012).

Several lines of evidence had suggested the effectiveness of PILI. A recent retrospective study investigating PILI's effects on LTs' language abilities found that LTs' communicative participation and expressive and receptive skills significantly increased following the intervention (Kwok et al. 2019). Concerning parents' communication skills following PILI, Sokmum et al. (2017) reported that parents in the PILI group showed an increase in using facilitative communication strategies during parent-child interaction, while the parents in the one-to-one speech therapy group remained consistent throughout the intervention processes. Besides, vocabulary, communication, and social skills were also improved in the PILI group (Sokmum et al. 2017).

A few reviews have also been conducted to examine the outcomes of PILI for children with language problems. Two meta-analysis studies highlighted that PILI had a significant, positive impact on young children's language abilities, both with and without cognitive issues (Heidlage et al. 2019; Roberts & Kaiser 2011). Moreover, Tosh et al. (2017), in their systematic review, reported that PILI was more effective in improving children's speech and language skills than no intervention. Although the studies mentioned earlier successfully confirmed PILI's effectiveness, there were a few limitations. The reviews included children with various speech and language problems, and the explanation about the effects of the structure and design of PILI programs was not offered. Available systematic review studies which focused on interventions for LTs (i.e., Cable & Domsch 2011; Deveney et al. 2017), on the other hand, discussed only different types of intervention approaches (i.e., PILI and clinician-directed intervention) and their effect on LTs' language skills.

Due to the gap mentioned above, factors related to the design of each PILI program that may affect the effectiveness of the programs are unidentified. Heidlage *et al.* (2019) argued that it was essential to examine and describe PILI programs' active ingredients to advance both research and practice. Furthermore, as there are two classifications of LTs available in the literature, it highlights the need to look in detail at the participants' characteristics

in PILI studies to aid the researchers' and clinicians' evaluation in choosing the appropriate PILI programs to be used.

The current work responds to the above needs by reviewing and discussing PILI programs' characteristics, including the design and structure, participants' criteria, and PILI programs' effectiveness. Therefore, the aims of the current scoping review are threefold. The first aim is to identify and describe the characteristics of participants in the reviewed studies. The second aim is to explore the design of the available PILI programs used explicitly in the intervention of LTs in terms of; a) intervention approaches, b) teaching strategies, c) duration of intervention, e) level of intervention, and f) intervention setting. The final aim is to explore the effectiveness of the PILI programs in relation to the characteristics of the studies.

## **METHOD**

The scoping review was performed by adhering to the framework proposed by Arksey and O'Malley (2005). This framework was selected as it offered a rigorous and transparent method throughout the reviewing process that could increase the reliability of the study outcomes. The stages of the framework were: 1) identifying research question(s), 2) identifying relevant studies, 3) study selection, 4) charting the data, and lastly, 5) collating, summarizing, and reporting the results.

## IDENTIFYING THE RESEARCH QUESTION(S)

The following questions guided this scoping review:

- 1. What are the characteristics of the participants in PILI studies?
- 2. What are the different approaches used in the available PILI programs?
- 3. What teaching strategies have been used in the available PILI programs?
- 4. What is the duration of the available PILI programs?
- 5. What are the levels of intervention used in the available PILI programs?
- 6. What intervention settings have been applied in the available PILI programs?
- 7. Are the available PILI programs effective in improving LTs' language abilities?

#### **IDENTIFYING RELEVANT STUDIES**

Arksey and O'Malley (2005) asserted the significance of being as comprehensive as possible in identifying relevant studies to be embedded in a scoping review. In order to achieve this, inclusion and exclusion criteria were listed.

#### INCLUSION AND EXCLUSION CRITERION

Prior to the identification of studies, the following inclusion and exclusion criteria were selected:

- 1. Article. As the study was conducted in 2018, only articles published between 1980 and 2018 and written in English or Malay language with full text available were selected. The time frame was considered appropriate as PILI gained popularity among professionals at the end of the 1970s (Mahoney et al. 1999). Unpublished dissertations, conference presentations, concept papers, and review articles, nonetheless, were omitted.
- 2. Participant. The child participants included in the studies were identified as late talking or having language delay or expressive language delay despite normal cognition, motor, sensory, and neurological systems, and aged between 18 and 42 months. Studies that included children with language problems associated with any known etiologies were discarded.
- 3. *Intervention*. Studies that used parents as the main interventionist were included. The parents must receive direct training or consultation from professionals. The PILI programs used were described in detail that they could potentially be replicated.

## SEARCH STRATEGY

The five databases selected for this review, namely ProQuest, Science Direct, Scopus, Taylor & Francis, and Wiley Online Library, were searched in July 2018. The following search terms were applied: parent\* AND intervention OR therap\* AND "late talk\*" OR "late language emergence" OR "early language delay" OR "expressive language delay". The reference lists of prior meta-analyses and systematic reviews (see Cable & Domsch 2011; Deveney *et al.* 2017; Roberts & Kaiser 2011; Tosh *et al.* 2017) were reviewed to identify studies

that met the inclusion criteria but were missed by the searches.

## STUDY SELECTION

Figure 1 illustrates the process of searching and reviewing the articles across the five selected databases. The search yielded 940 articles for review. The search results were then transferred into a Microsoft Excel document for comparison to identify duplicate articles. Following that, duplicate articles were excluded (n= 21). Next, the first and second reviewers independently selected the articles based on the inclusion criteria. Any emerging discrepancy during this process was discussed to reach a consensus. The first selection step was solely based on the titles of the articles listed in the earlier Microsoft Excel document. The titles were screened based on the inclusion and exclusion criteria. Therefore, the title that was clearly out of the scope of this study or mentioned the involvement of children with speech disorders and other developmental disorders and older children were removed. In addition, review articles also were discarded. A total of 60 articles were retained after this selection process. After that, the selection process involved abstract screening. The abstracts of the articles were collected in the same Excel document. The articles were then selected solely based on information from the abstract by again adhering to the inclusion and exclusion criteria. As a result, 38 articles were excluded as they did not fulfill the inclusion criteria. Hence, 22 relevant articles were retained and included in the full-text screening. An additional four articles from reference lists of past reviews, two each from Roberts and Kaiser (2011) and Tosh et al. (2016), were included in this final screening. Hence, 26 articles were screened from this, and only 15 studies were accepted for data extraction.

## CHARTING THE DATA

The first and second reviewers agreed upon a template created on Microsoft Excel for data charting purposes. Both reviewers independently charted relevant information on participant and intervention characteristics. The description of the charted information is shown in Table 1. Other data items that were considered during the charting process were; year, country of origin, and research design. These data items were incorporated as they gave vital information about the study. A meeting was convened to resolve disagreements by coming to a consensus.




Figure 1. Flow diagram illustrating the scoping review process

## Table 1. Charted information for the reviewed studies. Charted Information Participant' Child's participant Characteristics • Age • Gender • Types of language problems: expressive language delay or receptive and expressive language delay • Primary language Adult's participant • Gender • Ethnicity • Education level • Socioeconomic status Intervention' Intervention approaches Characteristics • Child-led: promote language learning in naturalistic contexts and responsive interaction • Adult-led: promote structured activities prepared by adults • Hybrid: promote language learning in naturalistic contexts, responsive interaction, and prompting strategies to encourage speech and language development. Teaching strategies • Strategies used to deliver the intervention. It may include lectures, demonstration (i.e., live, video), practical, discussion, and delivery of assignment

Duration of intervention

Duration of each training sessionTotal duration of intervention

to be continue....

Level of intervention

- · Individual
- Group

Intervention setting

- Clinic
- Home

Effectiveness of PILI programs

Design of the studies

- Groups used for comparison (e.g., PILI and no/delayed intervention, PILI and direct therapy,
- PILI, and other intervention)
- Results of the reviewed studies
- · Effect size

## COLLATING, SUMMARIZING, AND REPORTING RESULTS

The first and second reviewers scrutinized the 15 selected articles to answer the research questions. From these 15 articles, two studies were linked with several publications (see Buschmann et al. 2009, 2015; Girolametto et al. 1996a, 1997), while another article probed into two related studies that involved different participants (see Gibbard 1994). Although articles by Buschmann et al. (2009, 2015) involved different authors, the study in those articles had used the same participants and intervention program. This was a similar case with articles by Girolametto et al. (1996a, 1997). Thus, in order to ensure that the number of publications did not affect the review findings, the two publications by Buschmann et al. (2009, 2015) and Girolametto et al. (1996a, 1997) were considered as one each. Meanwhile, the article by Gibbard (1994) was treated as two studies.

#### **FINDINGS**

A total of 14 studies from five countries were identified for this review. Of these, five studies were conducted in the United States, five in the United Kingdom, two in Canada, and one in the Netherlands and Germany. Table 2 presents the summative descriptions of the 14 studies. In this section, the findings are presented based on the research questions outlined in Section 2. 1 and described in three sub-sections: (i) characteristics of participants; (ii) characteristics of PILI programs; and (iii) effectiveness of PILI programs.

## CHARACTERISTICS OF PARTICIPANTS

The families of LTs receiving PILI in the reviewed studies were culturally and economically diverse. A summary of participants' characteristics is also given in Table 2, and further explained in the following subsection.

#### CHILD PARTICIPANTS

Across the 14 selected studies, there were 519 child participants; 280 males and 102 females. Information about gender, however, was not provided for 137 children in five studies. In addition, two studies failed to report the exact age of the participating children (i.e., Girolametto *et al.* 1996b; McDade & McCartan 1998). Both studies only mentioned the inclusion criteria for child participants, which was between 24 and 42 months old. The mean age of the participating children in the remaining reviewed studies ranged from 26 to 35 months old at the beginning of the studies.

Of the 519 child participants, there were 455 LTs and 64 typical children. Eight studies only included children with expressive language delay, while five studies included children with either expressive language delay or receptive and expressive language delay, and one study included children with either language delay or speech and language delay. Typical children were only included in two studies as the control group. Although the diagnosis of the participating children differed from one study to another, all the studies incorporated similar inclusion criteria, whereby the children were at a specific age range, and their language problem was not secondary to other issues. As for language use, six studies focused on children with English as their primary language (i.e., Baxendale & Hesketh 2003; Girolametto et al. 1996b; Littleton Jr. 2004; Roberts et al. 2014; Roberts & Kaiser 2012), one study focused on German-speaking children (i.e., Buschmann et al. 2009), and the remaining seven studies did not include information regarding the language use of the participating children.

## ADULT PARTICIPANTS

Across the 13 studies, 205 adult participants were reported as directly involved in the intervention. Nonetheless, one study failed to report the number of adult participants

Table 2. Summative descriptions of the reviewed studies

		Table 2. Summative	table 2. Summative descriptions of the reviewed studies	wed studies	
Author (s)	Participants	Intervention	Outcome Measure (s)	Standardised Measure(s) Used	Results
Baxendale and Hesketh (2003)	Child  Population: Expressive language delay or receptive and expressive language delay  Age: 29-41 months  Gender: 7 females; 30 males  Primary Language: English  Adult  Ethnicity: Caucasian  School leaving age: 16 years (16), 18 years (1), further education (2)  Social/socioeconomic status: 'high level of social deprivation.	Design: Non-randomized controlled Program: Hanen Program Groups: intervention and direct therapy	Child: Receptive and expressive language skills, MLU Parent: Parent language- modeling techniques	Child: PLS-3-UK Parent: NA	Children in both groups improved their receptive and expressive abilities at post-test. The mean of the PLS-3 scores and MLU was higher in the intervention group than in the direct therapy group after 6 and 12-month, but the differences were not significant.  The mean of parent responses to child utterances for both groups increased after 6 months but decreased after 12 months—no significant difference between the groups in the percentage of parent responses at any assessment points.
Buschmann <i>et al.</i> (2009; 2015) n = 20 control	Child  Population: Typical language development or expressive language delay  Age: 24-27 months Gender: 39 females; 44 males Primary Language: German Adult Ethnicity: NR Education level: No graduation (3), middle school (9), high school (12) Social/socioeconomic status: NR	Design: Randomised controlled trial Program: Heidelberg Parent-Based Language Intervention Groups: intervention, no intervention, no intervention, anguage	Child: Receptive and expressive language skills (2009); working memory (2015) Parent: NA	Child: CDI-German, SETK (2009); AWST-R, K-ABC (2015) Parent: NA	The normal language group had significantly higher scores than the intervention and no intervention groups in all post-test language measures (p <0.05).  Children in the intervention group showed a significantly higher gain in both receptive and expressive language skills than children in the no intervention group with small to large effects (p<0.05; d=0.03-1.16) (2009).  Children in the intervention group had significantly better language comprehension (p<0.05; d=0.68), phonological memory (p<0.05; d=0.61-1.44), and episodic buffer than in the no intervention group (p<0.05; d=0.74) at post-test (2015).

•			
ς			
7	٠	٠	
	3	:	
5	٩	۶	
2	2	:	
Ç	1	:	

Gibbard (1994), Experiment 1	Child Population: No or little expressive language Age: 27-39 months Gender: 11 females, 25 males Primary Language: NR Adult Ethnicity: NR Education level: NR Social class: 1 (3), II (2), IIIM (8), IV (1), V (1), and unemployed (3)	Design: Randomised controlled trial Program: Parent Based Intervention Groups: intervention and no intervention	Child: Receptive and expressive language skills, MLU, grammatical ability, expressive vocabulary Parent: NA	Child: RDLS, Renfrew Action Picture Test, DLSPT Parent: NA	The mean score of expressive language skills for both groups improved at post-test. Children in the intervention group significantly showed greater gain than children in the no intervention group (p<0.01)—effect size n.s.
Gibbard (1994), Experiment 2	Child  Population: No or little expressive language  Age: 27-39 months  Gender: 6 females, 19 males  Primary Language: NR  Adult  Ethnicity: NR  Education level: NR  Social class: II (4), IIIM (4), IIIN  (4), IV (2), unemployed (2), and forces (1)	Design: Randomised controlled trial Program: Parent Based Intervention Groups: intervention 1, intervention 2 (control), direct therapy	Child: Receptive and expressive language skills, MLU, grammatical ability, expressive vocabulary Parent: NA	Child: RDLS, Renfrew Action Picture Test, DLSPT Parent: NA	There were differences in the expressive language gain between children in the intervention 1 group and the direct therapy group, with those from the intervention 1 group being higher. However, the differences were not significant.  The gain made (i.e., MLU, total score DLSPT) by the direct therapy group was significantly higher than those made by the intervention 2 (control) group (p<0.05)-effect size n.s.
Gibbard <i>et al.</i> (2004)	Child  Population: Language delay or speech and language delay  Age: 22-32 months  Gender: 5 females, 17 males  Language: NR  Total number: 12 parents  Adult  Ethnicity: NR  Education level: NR  Social class: II (3), IIIM (1), IV (4), V (1), unemployed (1), and armed forces (2)	Design: Non-randomized controlled trial Program: Parent Based Intervention (PBI) Groups: PBI and general care group	Child: Receptive, expressive language skills; expressive vocabulary, MLU Parent: NA	Child: RDLS; PLS-3 UK Parent: NA	The differences between the post-test scores of the intervention and general care group were significant (<0.05), with those in the intervention group showing higher scores in all language measures except for estimated vocabulary than in the direct therapy group—effect size n.s.

continuation					
Girolametto et al. (1996a)	Child Population: Expressive language delay Age of inclusion criteria: 24-42 months Gender: 5 females, 11 males Primary Language: English Adult Ethnicity: NR Education level: High school and post-secondary school Social/socioeconomic status: NR	Design: Randomised controlled trial Program: Hanen Program with focused stimulation Groups: Intervention and delayed intervention	Child: Expressive vocabulary (target words), play gesture, behavioral/emotional development Parent: NA	Child: CDI CBCL Parent: NA	Children in the intervention group significantly used more target words in a semi-structured task and increased use of symbolic play gestures than children in the delayed intervention group during post-test (p<0.03)—effect size n.s.  Parents in the intervention group reduced ratings for their children's aggressive/destructive behavior, whereas no rating reduction was observed among parents in the delayed treatment group. The differences were statistically significant (p<0.02)—effect size n.s.
Girolametto <i>et al.</i> (1996b; 1997)	Child  Population: Expressive language delay  Age: 23-33 months  Gender: NR  Primary Language: English  Adult  Ethnicity: Asian Canadian (1) and Caucasian (11)  Education level: High school (4) and post-secondary school (21)  Social/socioeconomic status:  Middle class	Design: Randomised controlled trial Program: Hanen Program with focused stimulation Groups: Intervention and delayed intervention	Child: Expressive vocabulary, 'talkativeness,' structural complexity (1996), consonant inventory, syllable structure levels, consonants correctly produced (1997) Parent: Parent: Parents' language - modeling techniques (1996)	Child: CDI (1996) Parent: NA	Mothers in the intervention group significantly (p<0.01) gave slower, less complicated, and used a greater number of target words and more focused stimulation than mothers in the delayed treatment group following intervention (1996). Effect size n.s. Children in intervention groups had larger vocabularies (p<0.02) and used more different words in interaction (p<0.01) than those in the delayed-intervention group at post-test. Effect size n.s.  Children in the intervention group exhibited more phonological diversity (p<0.05; d=0.6-1.1) following intervention than those in the delayed-intervention group, but there was no effect of treatment on the accuracy of speech production (1997)
Littleton Jr. (2004)	Child  Population: Expressive language delay  Age: 22-30 months  Gender: NR  Primary Language: English  Adult  Ethnicity: NR  Education level: Post-secondary education (8)  Social/socioeconomic status: NR	Design: Randomised controlled trial Program: Parent ACTION for Language Program (PALP) Groups: PALP and traditional early intervention	Child: Expressive vocabulary, talkativeness, and structural complexity.  Parent: Talkativeness, complexity of language input, and use of labels	Child: SICD, CDI Parent: NA	Children in the PALP group significantly increased their symbolic play skills to a greater degree than those in the traditional early intervention group (p<0.05). The gain in language measures was not significant at post-test between the groups.  The decrease in mean of parents' talkativeness and sentence complexity in both groups decreased following intervention were not significantly different between both groups.
					to be continue

	ntervention group (p≤0.01), whereas and for the children at size n.s.  nt on interactive al time for rvention group nges occurred in the n.s.	had greater gain a group on the PLS-4 ive communication (p=0.02; 0.75) ad a significantly es usage than parents 01; d=1.81-3.19).	orovement across all strated increases strategy after
	Children's language scores in the intervention group increased significantly at post-test (p≤0.01), whereas no significant differences were found for the children in the no-intervention group. Effect size n.s. Significantly increased of time spent on interactive engagement and decreased of actual time for unengagement (p≤0.05) in the intervention group following the intervention. No changes occurred in the no-intervention group—effect size n.s.	Children in the intervention group had greater gain than children in the no intervention group on the PLS-4 total (p=0.03; d=0.60) and expressive communication (p=0.04; d=0.67) score and TNW (p=0.02; 0.75) following the intervention.  Parents in the intervention group had a significantly higher percentage of EMT strategies usage than parents in the no-intervention group (p<0.01; d=1.81-3.19).	Three of four children showed improvement across all language measures. Parents demonstrated increases in their use of language facilitation strategy after instruction.
	Child: PLS-3, Engagement States Coding Scheme	Child: PLS-4 Parent: NA	Child: PLS- 4, EOWPVT Parent: NA
	Child: Receptive and expressive language skills, joint engagement	Child: Receptive and expressive language skills, NDW, MLUm, TNW Parent: Parents' use of language facilitation strategies	Child: Receptive and expressive language skills, NDW, MLUm, TNW Parent: Parents' use of language facilitation strategies
	Design: Non-randomized controlled trial Program: Hanen Program Groups: intervention and no intervention	Design: Randomized controlled trial Program: Enhanced Milieu Teaching (EMT) Groups: intervention, no intervention, normal language	Design: Single subject design Program: Enhanced Milieu Teaching Groups: NA
	Child Population: Expressive language delay Age: referred at 24 months Gender: NR Primary Language: NR Adult Ethnicity: NR Education level: NR Social/socioeconomic status: NR	Child  Population: Typical language development or receptive and expressive language delay Mean age: 29.86 – 31 months Gender: 9 females, 53 males Primary Language: English Adult Ethnicity: African American (3), Caucasian (12), other (1) Education level: College (2), Degree (6), Master (7), not respond (1) Social/socioeconomic status: Middle class	Child  Population: Receptive and expressive language delay  Age: 25-38 months  Gender: 2 females, 2 males  Primary Language: English  Adult  Ethnicity: White  Ethacation level: High School (1), college (2), Master (1)  Family annual income: USD50,000  -> USD100,000
continuation	McDade and McCartan (1998)	Roberts and Kaiser (2012)	Roberts et al. (2014)

Roberts and	Child	Design: Randomized	Child:	Child:	There were significant differences of all measures
Kaiser (2015)	Population: Receptive and	controlled trial	Receptive and	PLS- 4, CDI,	in receptive language (p=0.04; d=0.27-0.35) and every solve to a system of the system
	Age: 24-42 months  Gender: 18 females, 79 males	Teaching (EMT) Groups: EMT and no	skills, receptive and expressive	LO W1 V 1, 11 V 1	intervention between EMT and no intervention group
	Primary Language: NR Groups: 45 PILI, 52 usual-care	intervention	vocabulary, NDW		Parents in the intervention groups improved their use of all language facilitation strategies. The differences
	Adult Ethnisita: African American (9)		Parent:	Parent:	between the EMT and no intervention group were
	White (35), other (2)  Education level: High school (18), Undergraduate degree (17), Graduate degree (10) Family annual income: USD 60,000-		ratents use of language facilitation strategies	W.	significant (p.v.o.o.), p-v.45-5.19)
	0.00 / 1,000				
Van Balkom et	Child	Design: Randomized	Child:	Child:	Children in the PVHT group showed significant
al. (2010)	Population: Receptive and expressive language delay Age: 26-37 months	controlled trial  Program: Parent Based Video Home Training	MLU, grammaticality (syntax and	Dutch version of LARSP and RLDS	short-term and long-term improvement in grammar (p=0.015) and conversation coherence (p=0.008) as compared to children in the direct therapy group post-
	Gender: NR Primary Language: NR	(PVHT) Grouns: PVHT and direct	morphology),		intervention and after three months. Effect size n.s.
	Adult Ethnicity: NR	therapy	conversational coherence		
	Education level: NR			Parent:	
	social/socioeconomic stants: Middle class		rarent: NA	Y.	
Whitehurst et	Child	Design: Randomized	Child:	Child:	Children in the intervention group had a significantly
al. (1991)	Population: Expressive language disorder	controlled trial Program: Home-Based	Receptive and expressive	PPVT, ITPA, EOWPVT.	better score on the language measures than the no intervention group following intervention (p<0.05)—
	Mean age: 27.95 months	Intervention	vocabulary,	Templin-	effect size n.s.
	Gender: NR	Duration: 7 biweekly	expressive verbal	Darley Tests of	
	Language: NK Adult	sessions clinic visit (30 minufes/session)	nuency, articulation	Articulation	
	Ethnicity: NR	Groups: intervention and no		Parent:	
	Education level: NR	intervention	Parent:	NA	
	Social/socioeconomic status: NR		NA		

continuation...

Note: PLS: Pre-school Language Scales; CDI: Communicative Development Inventories; SETK: shandardised and norm-referenced instrument to examine the language status of German-speaking children; AWST-R: Expressive Vocabulary" Year-Old Children, Revision; K.-ABC: Kaufman Assessment Battery for Children; RDLS/RLDS: Reynell Developmental Language Scales; DLSPT: Derbyshire Language Scheme Picture Test; CBCL: Child Behavior Checklish; to -7 Test for Primary; REC: Revised Experiencer Constraint; PLAI: Preschool Language Assessment Instrument; SICD: Sequenced Inventory of Communicative Development; EOWPVT: Expressive One-Word \*-1-Fst of Language Development :\*\*TOLD Picture Vocabulary Test-Fourth Edition; MLUm: mean length of utterance in morphemes; NDW: number of different words; TNW: number of total words; PPVT: Peabody Picture Vocabulary Test; LARSP: Language Assessment, Remediation, "and Screening Procedure; ITPA: Illinois Test of Psycholinguistic Abilities; NR: not reported, NA: not applicable, n.s. not specified

involved in the treatment and no treatment groups (i.e., Whitehurst *et al.* 1991). Out of 205, 189 were mothers, three were fathers, and one was a grandmother. Information about gender was omitted for 12 adult participants in a study (i.e., Gibbard *et al.* 2004). Only five of the 14 selected studies gave information about the participants' ethnicities (i.e., Baxendale & Hesketh 2003; Girolametto *et al.* 1996a; Roberts & Kaiser 2015, 2012; Roberts *et al.* 2014). Among the included ethnicities were Caucasian/White (n=81), African American (n=11), Asian Canadian (n= 1), and others (n=3). As most of the adult participants were parents, the term 'parents' was used to describe and discuss the reviewed studies.

Six of 14 studies reported that most parents were high school graduates or higher (i.e., bachelor's and graduate degrees) (Girolametto et al. 1996a, 1996b; Littleton Jr. 2004; Roberts & Kaiser 2015, 2012; Roberts et al. 2014). On the contrary, most parents in two studies by Baxendale and Hesketh (2003) and Buschmann et al. (2009, 2015) did not complete high school education. As for annual family income, only two studies provided the information (i.e., Roberts & Kaiser 2015; Roberts et al. 2014). The reported family annual income in both studies ranged from an average of USD 50,000 to more than USD 100,000. Although three studies did not mention in detail the socioeconomic status of their participants, they claimed that either majority or all of the participants were from a middle-class family (Girolametto et al. 1996a; Roberts & Kaiser 2012; Van Balkom et al. 2010).

In general, most studies conducted before the 2000s gave incomplete demographic information, which hindered one from drawing a valid and definitive conclusion. Only three studies provided complete demographic information (i.e., Baxendale & Hesketh 2003; Roberts *et al.* 2014; Roberts & Kaiser 2012). Moreover, it was apparent from this limited demographic information that most mothers would join the intervention. Furthermore, although there were standard criteria to identify LTs, language problems faced by the child participants were still varied.

# CHARACTERISTICS OF PILI PROGRAMS

# CONTENT AND APPROACHES OF PILI PROGRAMS

There were eight PILI programs used across the studies. The programs were; (1) Hanen Program (HP), (2) adapted HP with Focused Language Stimulation (FLS), (3) Parent ACTION for Language Program (PALP), (4) Heidelberg Parent-Based Language Intervention (HPLI), (5) Parent-based Video Home Training (PVHT), (6) Enhanced Milieu Teaching (EMT), (7) Home-Based Treatment (HBT) and,

(8) Parent-Based Intervention (PBI). Most programs promoted teaching strategies in naturalistic contexts and responsive interaction and trained parents to adapt learned strategies in singing, shared games and reading books activities, except for the PBI. Moreover, other elements were included in the programs, such as discussing the types of play and children's play skills, dealing with media, and steps to improve children's auditory discrimination abilities and imitation skills. Interestingly, the PILI programs could also be further classified into three different approaches; i) child-led, ii) adult-led, and iii) hybrid approaches. A summary of the contents and intervention approaches for each PILI program is given in Table 3.

## CHILD-LED APPROACH

Seven studies employed the child-led approach in their PILI programs. The programs were HP (Baxendale & Hesketh 2003; McDade & McCartan 1998), FLS (Girolametto et al. 1996b, 1996a, 1997), PALP (Littleton Jr. 2004), PVHT (Van Balkom et al. 2010), and HPLI (Buschmann et al. 2009, 2015). These programs promoted parents to allow their children to lead the activities and trained the parents to use suitable language facilitation techniques while interacting with the children. The techniques included observing, waiting for a response, emphasizing keywords, and repeating and responding to the children's communication attempts. Moreover, the parents were taught to imply the techniques in their daily routine activities.

## ADULT-LED APPROACH

Amongst the 14 studies, three studies applied the adult-led approach in the PILI program; PBI (Gibbard 1994; Gibbard et al. 2004). The main objectives of PBI were to; i) increase a child's linguistic complexity from one-word level to three to four words per utterance; ii) improve a child's listening skills and auditory discrimination abilities; iii) increase a child's imitation skills; iv) educate parents on the importance of play, and v) develop the parents' ability and confidence to be the main therapist for their children (Gibbard 1998). In line with the concept of adult-led approach, whereby activities were selected by adults based on the child's interest to encourage learning, parents in this program were exposed to structured teaching situations and possible games to be conducted at home in order to meet the program's objectives. The parents were advised to encourage children to imitate (i.e., actions and/or speech) and use techniques, such as prompting, modelling, and forced alternatives, during the activities (Gibbard 1998).

Table 3. Content and duration of the parent-implemented language intervention (PIL1) programs for late talkers (LTs)

	Table 5. C	Table 3. Collient and duration of th	re parent-implemented tanguage intervention (f.1L.1) programs tot rate tarkers (f.1s)	guage miervention (FIL	1) programs for fate talk	ers (LIS)	
Hanen Program (HP)	Hanen Program + Focused language stimulation (FLS)	Parent ACTION for Language Program (PALP)	Heidelberg Parent- Based Language Intervention (HPLI)	Parent-based Video Home Training (PVHT)	Enhanced Milieu Teaching (EMT)	Home-Based Treatment (HBT)	Parent Based Intervention (PBI)
		Child-led approach			Hybrid a	Hybrid approach	Adult-led approach
Teaching strategies in naturalistic contexts & responsive interaction     action	Teaching strategies in naturalistic contexts & responsive interaction	Teaching strategies in naturalistic contexts & responsive interaction      Adaptation of	Normal language development and language delay     Teaching strategies in naturalistic con-	Strategies to create conversational coherence	Teaching strategies in naturalistic contexts & responsive interaction	Teaching strategies in naturalistic contexts & responsive interaction  2. Promoting strate-	The importance of play to child's language development     ment     Auditory discrimi- 2. Auditory discrimi-
<ol><li>Types of play and children's play skills</li></ol>	2. Types of play and children's play skills	learned strategies in singing, shared games, and reading activities	texts & responsive interaction 3. Adaptation of		<ul><li>2. Prompting strategies</li><li>3. Adaptation of</li></ul>	gies 3. Adaptation of learned strategies	nation abilities 3. Imitation skills
3. Adaptation of learned strategies in singing, shared games, and reading activities	3. Adaptation of learned strategies in singing, shared games, and reading activities  4. Frequently models elselected target words	3. Frequently model selected target words	learned strategies in singing and dancing, games, reading, and daily routine activities		learned strategies in reading, eating snacks, and daily routine activities	during story-time	4. Structured language teaching strategies
			DI	Duration			
11-12 weeks treatment	10-11 weeks treatment	12 weeks treatment	6 months treatment	13 weeks treatment	3 months treatment	7 biweekly sessions: 30 minutes	6 months treatment
- 8-9 group sessions: 2 hours 15 minutes - 3 home visits	- 7-8 group sessions: 2 hours 30 minutes - 3 home visits	- Once a week home visit: 60-90minutes	- / sessions within 3 months: 2 hours - 1 session 6 months later: 3 hours	- 6 home visits fortnightly: 90 minutes	- 4 workshops:  1 hour - 24 practice sessions twice per week: 40 -60 minutes		fortnightly: 60-75 minutes

Four studies had implemented the hybrid approach in their PILI programs, namely the EMT (Roberts & Kaiser 2015, 2012; Roberts *et al.* 2014) and HBT (Whitehurst *et al.* 1991). Both programs trained the parents to manipulate the children's natural environment and use milieu teaching procedures (i.e., model, mand-model, incidental teaching, and time delay) during their interactions. Response demands from children were not required during the procedure. Other responsive interaction strategies, such as expanding children's utterances and balancing verbal turntaking, were also emphasised in the programs. In the EMT, parents were required to practice all the learnt interaction strategies while eating snacks, reading books, and while performing common household activities during home visit.

## TEACHING STRATEGIES

Some teaching strategies that were used in the PILI programs in most of the reviewed studies comprised of giving lectures, demonstration either live or using videos and allocating time for the practical session (Baxendale &

Hesketh 2003; Buschmann et al. 2009; Gibbard 1994; Gibbard et al. 2004; Girolametto et al. 1996b, 1996a; McDade & McCartan 1998; Roberts & Kaiser 2015, 2012; Roberts et al. 2014; Van Balkom et al. 2010). During the practical session, the trainers asked the parents to practice in a group (role-play) or individually with their children. Corrective feedback was given to the parents immediately or at the end of the practical session. Some training sessions embedded discussion and delivery of assignments or homework (Baxendale & Hesketh 2003; Girolametto et al. 1996b, 1996a, 1997; Littleton Jr. 2004; McDade & McCartan 1998; Whitehurst et al. 1991). The elaborated teaching strategies were not limited to group training, as individual training was incorporated as well. The design and structure of each PILI program can be seen in Table 4.

#### **DURATION OF INTERVENTION**

Various durations were noted for the PILI programs across the reviewed studies. Most programs were conducted for 11-12 weeks (Baxendale & Hesketh 2003; Girolametto *et* 

Table 4. Characteristics of the available parent-implemented language intervention (PILI) programs for late talkers (LTs)

			Teachin	g Stra	tegies			vel of vention	Interv Sett		_
	Program	Lecture	Video /Live Demonstration	Practical	Discussion	Homework	Individual	Group	Clinic	Home	Effectiveness
Child-led Approach	Hanen Program (HP)	/	/	/	/	/	/	/	/	/	Increased children's language scores and their engagement in the intervention group. No changes were observed in the no intervention group (McDade & McCartan, 1998). When compared with the direct therapy group, no significant difference in children language abilities and percentage of parents' responses can be observed (Baxendale & Hesketh, 2003)

to be continue...

continuation..

	Hanen Program + Focused language stimulation (FLS)	/	/	/	/	/	1	/	/	/	Children in the FLS grousignificantly had better expressive skills and phonological skills and were rated lower for their aggressive behavior than those in the delayed intervention group (Girolametto et al., 1996) 1996a, 1997)
											Mothers in the FLS group used more effective communication strategies than mothers in the delay intervention group (Girolametto et al., 1996)
	Parent ACTION for Language Program (PALP)		/	/	/	/	/			/	The PALP group significantly increased their symbolic play skills than those in the traditional earnetervention group—no significant difference in mothers' communicative behaviors were observed between the groups (Littleton Jr., 2004).
	Heidelberg Parent-Based Language Intervention (HPLI)	/	/	/		/		/	/		Significantly higher gain the intervention group in language measures (Buschmann et al., 2009) and better phonological memory and episodic buffer than the no interve- tion group (Buschmann et al., 2015)
	Parent-based Video Home Training (PVHT)		/	/	/		/			/	The PVHT group significantly showed short-tern and long-term improvement in grammar and conversation coherence (Van Balkom et al., 2010)
Adult-led Approach	Parent Based Intervention (PBI)	/	1	/	/			1	/		The PBI group significant showed greater gain that the no intervention group language measures. The differences in the gains between the PBI and direct therapy group were not significant (Gibbard, 1995 Study in 2004 revealed to The PBI group had highescores in language measures than the direct therapy group (Gibbard al., 2004)

continuation...

Hybrid Approach	Enhanced Milieu Teaching (EMT)	/	/	/	/		/	/	/	Children in the EMT showed improvement (Roberts et al., 2014) and had a greater gain in language measures than in the control group. Parents used on EMT strategies also increased significantly(Roberts & Kaiser, 2015; Roberts et al., 2014; Roberts & Kaiser, 2012)
	Home-Based Treatment (HBT)			/	/	/	/	/		Those in the HBT group had a significantly better score on the language measures than the no intervention group (Whitehurst et al., 1991)

al. 1996b, 1996a; Littleton Jr. 2004; McDade & McCartan 1998; Roberts & Kaiser 2015, 2012; Roberts et al. 2014; Van Balkom et al. 2010; Whitehurst et al. 1991), either once or twice a week or fortnightly. The longest programs were PBI (Gibbard 1994; Gibbard et al. 2004) and HPLI (Buschmann et al. 2009) that ran for six months. The duration for each individual session was between 30 to 90 minutes, with the shortest individual session in the HBT (Whitehurst et al. 1991) and the longest was in the PALP and PVHT (Littleton Jr. 2004; Van Balkom et al. 2010). As for the group session, the average duration was two hours per session. The shortest duration was one hour (Gibbard 1994; Gibbard et al. 2004), whereas the longest duration was three hours (Buschmann et al. 2009) for the group session. The EMT has the most number of training sessions and is conducted most frequently.

## LEVEL OF INTERVENTION

From the review, it was noted that the interventions were performed as either individual or group training. Six studies conducted individual training only (Littleton 2004; Roberts & Kaiser 2015, 2012; Roberts *et al.* 2014; Van Balkom *et al.* 2010; Whitehurst *et al.* 1991), four studies conducted group training only (Buschmann *et al.* 2009; Gibbard 1994; Gibbard *et al.* 2004), and another four conducted both group and individual training (Baxendale & Hesketh 2003; Girolametto et al. 1996b, 1996a; McDade & McCartan 1998).

The number of parents per training group, nevertheless, was not provided in most studies. Baxendale and Hesketh (2003) only mentioned that 19 late-talking children and their families took part in one of the five HP groups assessed in the study, whereas Girolametto *et al.* (1996b, 1996a)

asserted that the HP was administered to eight and 12 families, respectively, by two experienced SLPs. Although Buschmann *et al.* (2009) failed to specify the number of parents per training group in their study, their HPLI was designed for a group of five to 10 people. Gibbard *et al.* (2004), on the other hand, claimed that eight was the maximum number of parents per training group in their study as it was manageable in practice. In Gibbard's study, the parents were divided into two-parent training groups composed of eight and four parents.

## INTERVENTION SETTING

Three of the PILI programs in the reviewed studies were conducted at clinics and required home visits (Baxendale & Hesketh 2003; Girolametto et al. 1996b, 1996a; McDade & McCartan 1998; Roberts & Kaiser 2015; 2012; Roberts et al. 2014). During the clinic visit, the trainers focused more on delivering new information, teaching new techniques or strategies, facilitating discussion, and holding practicals. During the home visit, the trainers emphasized helping the parents apply the learned techniques at home with their children. For programs conducted in a clinic setting only (n=3), the participating parents were exposed to lectures, videos, role play, and discussion (Buschmann et al. 2009; Gibbard 1994; Gibbard et al. 2004; Whitehurst et al. 1991). The same exposure (i.e., direct teaching, show and discuss video) was given to parents who participated in the program conducted in a home setting only (n=2) (Littleton 2004; Van Balkom et al. 2010).

## EFFECTIVENESS OF PILI PROGRAMS

Most of the reviewed studies evaluated the effectiveness of the PILI programs by comparing language measures of children in different groups, either 1) PILI group and no / delayed intervention group or 2) PILI group and direct therapy group or 4) PILI group and other intervention. One study, however, applied a single subject design (i.e., Roberts et al. 2014). Regardless of the types of language problems, there were consistent findings across studies (n=8). LTs' language measures were significantly higher in the PILI group than the no/delayed intervention group with small to large effect sizes (i.e., Buschmann et al. 2009, 2015; Gibbard 1994; Girolametto et al. 1996a, 1996b, 1997; McDade & McCartan 1998; Roberts & Kaiser 2015; Roberts & Kaiser 2012; Whitehurst et al. 1991). Out of these eight studies, only two studies involved children with receptive and expressive language delay. These studies used the EMT, the individual basis PILI program, as their intervention. Other studies only involved children with expressive delay and used either group or individual and group (mixed) training.

When PILI was compared with direct therapy (n=3), mixed findings were found. Two studies found no significant difference between these groups following intervention (i.e., Baxendale & Hesketh 2003; Gibbard 1994). On the other hand, one study found that LTs' in the PILI groups had higher scores in grammatically correct utterances and conversation coherence than LTs' in the direct therapy group (i.e., Van Balkom et al. 2010). These findings suggest that PILI can be as effective as direct therapy. Similarly, findings from comparing the PILI and other intervention groups (n=2) were also varied. One study found superior effects of the PILI program (i.e., Gibbard et al. 2004), whereas another study found no significant effects of PILI from the traditional early intervention (non-SLP) (i.e., Littleton Jr. 2004). Although the control group in the study conducted by Gibbard (2004) had opportunities to meet speech therapists, they only received general verbal advice on language stimulation techniques to facilitate children's language development for two sessions. Thus, in the study, the PILI group was not compared with the direct therapy group; rather, the PILI group was compared with the 'other intervention' group. Out of eight PILI programs, only the PBI was conducted among children with only expressive delay (Gibbard 1994) and receptive and expressive delay (Gibbard et al. 2004). Other PILI programs, however, were evaluated only in one of the populations mentioned.

Interestingly only six of the reviewed studies included parents' communicative behaviors as their dependent variables. A similar pattern could be observed as studies comparing differences between parents in the PILI group and no/delay intervention group found a higher percentage of use of learned strategies in the PILI groups than the no/delay intervention group with medium to large effect sizes (i.e., Girolametto et al. 1996b; Roberts & Kaiser 2015; Roberts & Kaiser 2012). However, when compared with

the direct therapy or other intervention groups, no significant differences were found, although the mean of targeted communicative behaviors increased following the intervention (i.e., Baxendale & Hesketh 2003; Littleton Jr. 2004)

## **DISCUSSION**

The purposes of this scoping review study were to explore the studies' characteristics (i.e., participants, PILI programs) and describe the effectiveness of the PILI programs in relation to the characteristics. A systematic search of the literature from 1980 to 2018 yielded only 15 articles that fulfilled the predetermined inclusion criteria. Within these studies, eight different PILI programs were identified, and the characteristics of participants were varied. Six of the studies measured outcomes from both parents and children, while another half only measured the outcome of the children. Findings related to the effectiveness of the PILI programs also differed depending on the study design, type of PILI programs used, and involved participants.

The majority of the adult participants in all the reviewed studies were mothers. However, it was not clear whether the studies intended to involve only mothers during intervention or fathers themselves did not want to participate. This is noteworthy, as previous studies have shown many differences between mothers' and fathers' communication styles with their children. For instance, Leaper et al. (1998) found that fathers tended to talk less but used more directive and informing language than the mothers did. These differences occurred more likely among parents of younger children than for older children. Although fathers did not talk as much as mothers, their type-token ratio was denser than mothers (Kwon et al. 2013). These differences in communication style have been proven to have different effects on children's language development (Teufl et al. 2020). However, whether involving fathers during the intervention can replicate the effectiveness of the PILI programs is unknown.

Other than that, the child participants' language problems were varied, despite their criteria satisfying the characteristics of LTs. Some studies included children with both receptive and expressive language delay, whereas others included those with expressive language delay. The evaluation of the PILI programs interestingly showed that regardless of the differences in the language problems, the PILI programs were proven effective in improving children's language abilities and changing parents' communicative behaviors when comparisons were made between PILI and no/delayed intervention groups. The

findings were supported by other reviews that highlighted the positive effects of PILI on children's receptive and expressive language and parents' use of language facilitation strategies (e.g., Heidlage et al. 2019; Roberts et al. 2019).

Nevertheless, caution needs to be made in interpreting the findings. This is because studies involving children with receptive and expressive delay mainly implemented the EMT that consists of intensive individual training. A previous meta-analysis study revealed that individual parent training more benefited parents and children than group parent training (Lundahl et al. 2006). In addition, the larger intervention dosage of the EMT (measured by the total length of time receiving training) compared to other PILI programs might contribute to the effectiveness of the program among children with receptive and expressive delay (e.g., Maher et al. 2011). Interestingly, the effectiveness of other individual PILI programs was also observed in the PVHT, where the study that used the PVHT was the only study that found better effects of PILI than direct therapy conducted by clinicians. Although only half of the language measures were significant, the findings of the study were still promising. However, as individual PILI programs are usually conducted at home, they may cause higher costs and more time for the service provider (French & Yates 2018; Tosh et al. 2016).

Another important finding is that except for the PBI, other PILI programs were evaluated among children with either receptive and language delay or expressive delay only. Results of the studies indicate that the scores in language measures of children in the PBI group increased following the intervention (Gibbard 1994; Gibbard et al. 2004). However, the finding is just a piece of preliminary evidence, and a definite conclusion about the effectiveness of the PBI in both populations cannot be made. This is because only two studies were conducted among children with expressive delay, and one small study was conducted among children with both receptive and expressive delay using the PBI. Therefore, more studies are needed to evaluate whether the same PILI programs have similar effects on different child populations.

## CONCLUSION

This scoping review examined the literature on PILI studies by narrowing its focus on LTs. Viewing the results altogether, it is apparent that participants of PILI studies are diverse, and the structure and design of the PILI programs seemed varied across studies. Current evidence indicates that PILI is more effective than no/delayed intervention and can be as effective as direct therapy provided by clinicians. It is hoped that researchers and clinicians can use findings from this study to evaluate and choose the appropriate PILI programs that are most suitable and cater to the parents of late-talking children in their settings. Moreover, the findings are also hoped to be useful in developing new programs, especially in countries where evidence-based parental training programs are not yet available.

## **LIMITATIONS**

Although this scoping review was conducted by adhering to the framework prescribed by Arksey and O'Malley (2005), some noteworthy limitations must be highlighted. First, only English-language articles were reviewed in this study. The analysis omitted potentially relevant articles written in other languages. Secondly, the researchers in this study limited the searches to cover published articles only. Hence, any PILI program available in the grey literature was discarded during the review. Thirdly, only databases subscribed by the International Islamic University Malaysia (IIUM) were included. Fourth, it is possible that due to the selected terms used during searches, some articles on this topic were not identified, as the researchers might have overlooked some terms related to the study. Finally, there was also a risk of removing paper prematurely during the study selection as the first step involved screening articles solely based on titles.

#### **ACKNOWLEDGMENTS**

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article. This paper was supported by Fundamental Research Grant Scheme, Ministry of Higher Education Malaysia (FRGS/1/2019/SKK06/UIAM/03/1).

## **REFERENCES**

American Speech-Language-Hearing Association. 2008. Roles and responsibilities of speech-language pathologists in early intervention: guidelines [Guidelines]. doi:10.1044/policy.GL2008-00293

Arksey, H. & O'Malley, L. 2005. Scoping studies: Towards a methodological framework. *International Journal of Social Research Methodology: Theory and Practice* 8(1): 19–32. doi:10.1080/1364557032000119616

Armstrong, R., Arnott, W., Copland, D. A., Mcmahon, K., Khan, A., Najman, J. M. & Scott, J. G. 2016. Change in receptive vocabulary from childhood to adulthood: associated mental health, education,

- and employment outcomes. *International Journal of Language and Communication Disorders* 1–12. doi:10.1111/1460-6984.12301
- Baxendale, J. & Hesketh, A. 2003. Comparison of the effectiveness of the Hanen Parent Programme and traditional clinic therapy. *International Journal of Language and Communication Disorders* 38(4): 397–415. doi:10.1080/1368282031000121651
- Buschmann, A., Jooss, B., Rupp, A., Feldhusen, F., Pietz, J. & Philippi, H. 2009. Parent based language intervention for 2-year-old children with specific expressive language delay: A randomised controlled trial. Archives of Disease in Childhood 94(2): 110– 116. doi:10.1136/adc.2008.141572
- Buschmann, A., Multhauf, B., Hasselhorn, M. & Pietz, J. 2015. Long-Term Effects of a Parent-Based Language Intervention on Language Outcomes and Working Memory for Late-Talking Toddlers. *Journal of Early Intervention* 37(3): 175–189. doi:10.1177/1053815115609384
- Cable, A. L. & Domsch, C. 2011. Systematic review of the literature on the treatment of children with late language emergence. *International Journal of Language and Communication Disorders* 46(2): 138–154. doi:10.3109/13682822.2010.487883
- Dale, P. S., McMilan, A. J., Hayiou-Thomas, M. E. & Plomin, R. 2014. Illusory Recovery: Are Recovered Children With Early Language Delay at Continuing Elevated Risk? *American Journal of Speech-Language Pathology* 23(August): 437–447. doi:10.1044/2014
- Desmarais, C., Sylvestre, A., Meyer, F., Bairati, I. & Rouleau, N. 2008. Systematic review of the literature on characteristics of late-talking toddlers. *International Journal of Language & Communication Disorders* 43(4): 361–389. doi:10.1080/13682820701546854
- Deveney, S. L., Hagaman, J. L. & Bjornsen, A. L. 2017.
  Parent-Implemented Versus Clinician-Directed
  Interventions for Late-Talking Toddlers: A Systematic
  Review of the Literature. *Communication Disorders*Quarterly 1–10. doi:10.1177/1525740117705116
- Ebbels, S. H., McCartney, E., Slonims, V., Dockrell, J. E. & Norbury, C. F. 2019. Evidence-based pathways to intervention for children with language disorders. *International Journal of Language and Communication Disorders* 54(1): 3–19. doi:10.1111/1460-6984.12387
- Espe-sherwindt, M. & Serrano, A. 2016. It takes two: The role of family-centered practices in communication intervention. *Revista de Logopedia, Foniatría y Audiología*. doi:10.1016/j.rlfa.2016.07.006
- French, A. N. & Yates, B. T. 2018. Cost-Effectiveness of Parent-Child Interaction Therapy in Clinics versus Homes: Client, Provider, Administrator, and Overall Perspectives. *Journal of Child and Family Studies* (August). doi:10.1007/s10826-018-1159-4

- Gibbard, D. 1994. Parental-based intervention with preschool language-delayed children. *International Journal of Language & Communication Disorders* 29(2): 131–150. doi:10.3109/13682829409041488
- Gibbard, D. 1998. Parent-Based Intervention Programme: A Group Approach For Language-Delayed Children. Oxford: Winslow.
- Gibbard, D., Coglan, L. & MacDonald, J. 2004. Costeffectiveness analysis of current practice and parent intervention for children under 3 years presenting with expressive language delay. *International Journal of Language and Communication Disorders* 39(2): 229–244. doi:10.1080/136828203100016188 39
- Girolametto, L., Pearce, P. S. & Weitzman, E. 1996a. Interactive Focused Stimulation for Toddlers With Expressive Vocabulary Delays. *Journal of Speech Language and Hearing Research* 39(6): 1274–1283. doi:10.1044/jshr.3906.1274
- Girolametto, L., Pearce, P. S. & Weitzman, E. 1996b.
  The Effects of Focused Stimulation for Promoting Vocabulary in Young Children with Delays: A Pilot Study. *Journal of Children's Communication Development* 17(2): 39–49.
- Girolametto, L., Pearce, P. S. & Weitzman, E. 1997. Effects of Lexical Intervention on the Phonology of Late Talkers. *Journal of Speech Language and Hearing Research* 40: 338–348.
- Girolametto, L., Wiigs, M., Smyth, R., Weitzman, E. & Pearce, P. S. 2001. Children With a History of Expressive Vocabulary Delay. *American Journal of Speech-Language Pathology* 10(4): 358–369. doi:10.1044/1058-0360(2001/030)
- Hawa, V. V. & Spanoudis, G. 2014. Toddlers with delayed expressive language: An overview of the characteristics, risk factors, and language outcomes. *Research in Developmental Disabilities* 35(2): 400–407. doi:10.1016/j.ridd.2013.10.027
- Heidlage, J. K., Cunningham, J. E., Kaiser, A. P., Trivette,
  C. M., Barton, E. E., Frey, J. R. & Roberts, M. Y.
  2019. The effects of parent-implemented language interventions on child linguistic outcomes: A meta-analysis. *Early Childhood Research Quarterly* 50: 6–23. doi:10.1016/j.ecresq.2018.12.006
- Henrichs, J., Rescorla, L., Schenk, J. J., Schmidt,
  H. G., Jaddoe, V. W. V, Hofman, A. & Raat, H.
  2011. Examining Continuity of Early Expressive
  Vocabulary Development: The Generation R Study.
  Journal of Speech, Language, and Hearing Research
  54: 854–869. doi:10.1044/1092-4388(2010/09-0255)854
- Horwitz, S. M. C., Irwin, J. R., Briggs-Gowan, M. J., Bosson Heenan, J. M., Mendoza, J. & Carter, A. S. 2003. Language delay in a community cohort of young children. *Journal of the American Academy* of Child and Adolescent Psychiatry 42(8): 932–940. doi:10.1097/01.CHI.0000046889.27264.5E

- Kaderavek, J. N. 2011. Language disorders in children: Fundamental concepts of assessment and intervention. Pearson/Prentice.
- Kwok, E. Y. L., Cunningham, B. J. & Cardy, J. O. 2019. Effectiveness of a Parent-Implemented Language Intervention for Late-to-Talk Children: A Real-World Retrospective Clinical Chart. *International Journal* of Speech-Language Pathology 22(1): 48–58.
- Kwon, K., Bingham, G., Lewsader, J., Jeon, H.-J. & Elicker, J. 2013. Structured Task Versus Free Play: The Influence of Social Context on Parenting Quality, Toddlers 'Engagement with Parents and Play Behaviors, and Parent Toddler Language Use. *Child Youth Care Forum* 42: 207–224. doi:10.1007/s10566-013-9198-x
- Law, J, Dennis, J. A. & Charlton, J. J. 2017. Speech and language therapy interventions for children with primary speech and/or language disorders. *Cochrane Database of Systematic Reviews* (1). doi:10.1002/14651858.CD012490.www.cochranelibrary.com
- Law, James, Levickis, P., Rodríguez-Ortiz, I. R., Matić, A., Lyons, R., Messarra, C., Kouba Hreich, E., et al. 2019. Working with the parents and families of children with developmental language disorders: An international perspective. *Journal of Communication Disorders* 82(105922). doi:10.1016/j.jcomdis.2019.105922
- Leaper, C., Anderson, K. J. & Sanders, P. 1998. Moderators of Gender Effects on Parents 'Talk to Their Children: A Meta-Analysis. *Developmental Psychology* 34(1): 3–27.
- Leonard, L. B. 2014. Children with Specific Language Impairment. MIT press.
- Littleton Jr., R. F. 2004. The modifiability of language input with toddlers with expressive language delay: A study of a team approach to parent training. *ProQuest Dissertations and Theses* 197.
- Lundahl, B., Risser, H. J. & Lovejoy, M. C. 2006. A metaanalysis of parent training: Moderators and followup effects. *Clinical Psychology Review 26* 26: 86– 104. doi:10.1016/j.cpr.2005.07.004
- Maher, E. J., Marcynyszyn, L. A., Corwin, T. W. & Hodnett, R. 2011. Dosage matters: The relationship between participation in the Nurturing Parenting Program for infants, toddlers, and preschoolers and subsequent child maltreatment. *Children and Youth Services Review* 33: 1426–1434.
- Mahoney, G., Kaiser, A., Girolametto, L., MacDonald, J.,
  Robinson, C., Safford, P. & Spiker, D. 1999. Parent
  Education in Early Intervention: A Call for a Renewed
  Focus. *Topics in Early Childhood Special Education*19(3): 131–140. doi:10.1177/027112149901900301

- McDade, A. & McCartan, P. 1998. Partnership with parents: a pilot project. *International Journal of Language and Communication Disorders* 33(Supplement): 556–561. doi:10.3109/13682829809179485
- Paul, R. 1996. Clinical Implications of the Natural History of Slow Expressive Language Development. American Journal of Speech-Language Pathology 5: 5–21.
- Paul, R. 2001. Speech and language impairments in children. Causes, characteristics, intervention and outcome. (D. Bishop & L. Leonard, Eds.). East Sussex, UK: Psychology Press.
- Pring, T., Flood, E., Dodd, B. & Joffe, V. 2012. The working practices and clinical experiences of paediatric speech and language therapists: A national UK survey. *International Journal of Language and Communication Disorders* 47(6): 696–708. doi:10.1111/j.1460-6984.2012.00177.x
- Rescorla, L. 1989. The Language Development Survey: A Screening for Delayed Language in Toddlers. *Journal of Speech and Hearing Disorders* 54(November): 587–599.
- Rescorla, L. 2009. Age 17 Language and Reading Outcomes in Late-Talking Toddlers: Support for a Dimensional Perspective on Language Delay. *Journal of Speech, Language, and Hearing Research* 52(February): 16–30.
- Roberts, M. Y., Curtis, P. R., Sone, B. J. & Hampton, L. H. 2019. Association of Parent Training with Child Language Development: A Systematic Review and Meta-Analysis. *JAMA Pediatrics* 173(7): 671–680. doi:https://dx.doi.org/10.1001/jamapediatrics.2019.1197
- Roberts, M. Y. & Kaiser, A. P. 2015. Early Intervention for Toddlers With Language Delays: A Randomized Controlled Trial. *Pediatrics* 135(4): 686–693. doi:10.1542/peds.2014-2134
- Roberts, M. Y. & Kaiser, A. P. 2012. Assessing the Effects of a Parent-Implemented Language Intervention for Children With Language Impairments Using Empirical Benchmarks: A Pilot Study. *Journal of Speech Language and Hearing Research* 55(6): 1655–1670. doi:10.1044/1092-4388(2012/11-0236)
- Roberts, M. Y., Kaiser, A. P., Wolfe, C. E., Bryant, J. D. & Spidalieri, A. M. 2014. The Effects of the Teach-Model-Coach-Review Instructional Approach on Caregiver Use of Language Support Strategies and Children's Expressive Language Skills. *Journal of Speech, Language, and Hearing Research* 57(1): 1851–1869. doi:10.1044/2014
- Roberts, M. Y & Kaiser, A. P. 2011. The Effectiveness of Parent-Implemented Language Interventions: A Meta-Analysis. *American Journal of*

- *Speech-Language Pathology* 20(3): 180–199. doi:10.1044/1058-0360(2011/10-0055)amounts
- Scarborough, H. S. & Dobrich, W. 1990. Development of Children with Early Language Delay. *Journal of Speech and Hearing Research* 33(March): 70–83.
- Singleton, N. C. 2018. Late Talkers: Why the Waitand-See Approach Is Outdated. *Pediatric Clinics of North America* 65(1): 13–29. doi:10.1016/j. pcl.2017.08.018
- Sokmum, S., Joginder Singh, S. & Vandort, S. 2017. The impact of Hanen More Than Words Programme on Parents of children with ASD in Malaysia. *Jurnal Sains Kesihatan Malaysia* 15(2): 43–51. Retrieved from http://ejournals.ukm.my/jskm/article/view/14882/6221
- St Clair, M. C., Forrest, C. L., Kok Yew, S. G. & Gibson, J. L. 2019. Early Risk Factors and Emotional Difficulties in Children at Risk of Developmental Language Disorder: A Population Cohort Study. *Journal of Speech, Language, and Hearing Research* 1–22
- Teufl, L., Deichmann, F., Supper, B. & Ahnert, L. 2020. How fathers 'attachment security and education contribute to early child language skills above and beyond mothers: parent-child conversation under scrutiny. Attachment & Human Development 22(1): 71–84. doi:10.1080/14616734.2019.1589063
- Tosh, R., Arnott, W. & Scarinci, N. 2016. Parentimplemented home therapy programmes for speech and language: A systematic review. *International Journal of Language and Communication Disorders*. doi:10.1111/1460-6984.12280
- Van Balkom, H., Verhoeven, L., Van Weerdenburg, M. & Stoep, J. 2010. Effects of Parent-based Video Home Training in children with developmental language delay. *Child Language Teaching and Therapy* 26(3): 221–237. doi:10.1177/0265659009349978
- Whitehurst, G. J., Fischel, J. E., Lonigan, C. J., Valdez-Menchaca, M. ., Arnold, D. S. & Smith, M. 1991. Treatment of Early Expressive Language Delay: If, When, and How. *Topics in Language Disorders*.
- Zubrick, S. R., Taylor, C. L., Rice, M. L. & Slegers, D. W. 2007. Late Language Emergence at 24 Months: An Epidemiological Study of Prevalence, Predictors, and Covariates. *Journal of Speech, Language, and Hearing Research* 50(December): 1562–1592.

- <sup>1</sup>Nur Hanisah Tukiran, <sup>1</sup>Nor Azrita Mohamed Zain\*, <sup>1</sup>Natrah Ahmad Nordin, <sup>2</sup>Nadzirah Ahmad Basri
- 1Department of Audiology Speech-Language and Pathology, Kulliyyah of Allied Health Sciences, International Islamic University Malaysia, Jalan Sultan Ahmad Shah, 25000 Kuantan, Pahang, Malaysia
- <sup>2</sup>Department of Psychiatry, Kulliyyah of Medicine, International Islamic University Malaysia, Pahang, Malaysia.

### \*Corresponding author:

Department of Audiology and Speech-Language Pathology, Kulliyyah of Allied Health Sciences, International Islamic University Malaysia, Jalan Sultan Ahmad Shah, 25000 Kuantan, Pahang, Malaysia. Email: znazrita@iium.edu.my