

PATHOGENICITY OF ENTOMOPATHOGENIC FUNGI *Lecanicillium lecanii* AND *Beauveria bassiana* AGAINST *Pseudococcus jackbeardsleyi* (PSEUDOCOCCIDAE) INFECTING RAMBUTAN

Sempurna Ginting^{1*}, Djamilah², Tunjung pamekas², Hendri Bustaman², Priyatningsih², Sipriyadi³ & Risky Hadi Wibowo³

^{1,2}Plant Protection Department,
Faculty of Agriculture,
Universitas Bengkulu, Indonesia.

³Biology Department,
Faculty of Mathematics and Natural Sciences,
Universitas Bengkulu, Indonesia.

*Corresponding email: *sempurnaginting@unib.ac.id*

ABSTRACT

Pseudococcus jackbeardsleyi is an important pest that attacks rambutan fruit (*Nephelium lappaceum* L.). The *P. jackbeardsleyi* can cause severe damage in rambutan because it is able to survive in postharvest commodities. Therefore, is a need to control this pest to avoid losses. The species of *P. jackbeardsleyi* can be controlled by either chemical means using insecticides or biological means using entomopathogenic fungi. Between the two control measures, the use of entomopathogenic fungi is an environmental friendly control measure and causes no harm to human health. The purpose of this study was to examine the pathogenicity of entomopathogenic fungi *Lecanicillium lecanii* and *Beauveria bassiana* against *P. jackbeardsleyi* (Pseudococcidae) on rambutan. The application method was by spraying *P. jackbeardsleyi* with fungi conidia suspension. The data obtained was analysed using the Statistical Analysis System (SAS) program. The results showed that *L. lecanii* + soap was the most effective cause of mortality for *P. jackbeardsleyi*, then *L. lecanii*, and *B. bassiana* + soap, while the lowest was *B. bassiana*. Based on this study, the use of entomopathogenic fungi is effective against *P. jackbeardsleyi* in rambutan.

Keywords: Pathogenicity, *Lecanicillium lecanii*, *Beauveria bassiana*, *Pseudococcus jackbeardsleyi*, rambutan.

ABSTRAK

Pseudococcus jackbeardsleyi merupakan perosak penting yang menyerang buah rambutan (*Nephelium lappaceum* L.). Spesies *P. jackbeardsleyi* ini mampu menyebabkan kerosakan serius pada buah rambutan kerana spesies ini dapat bertahan pada peringkat komoditi lepas tuai. Oleh itu, spesies ini perlu harus dikawal agar tidak menyebabkan kerugian. Kawalan *P. jackbeardsleyi* dapat dilakukan secara kimia dengan menggunakan inseksitid ataupun dengan kawalan secara biologi menggunakan kulat entomopatogenik. Di antara kedua kawalan tersebut, penggunaan kulat entomopatogenik merupakan kawalan yang mesra alam dan tidak memberikan kesan

negatif terhadap kesehatan manusia. Kajian ini dijalankan adalah bertujuan untuk menguji kesan kepatogenan kulat entomopatogenik *Lecanicillium lecanii* dan *Beauveria bassiana* terhadap *P. jackbeardsleyi* pada rambutan. Kaedah yang digunakan dalam kajian ini adalah secara semburan *P. jackbeardsleyi* dengan campuran konidia. Data yang diperoleh dianalisis menggunakan perisian *Statistical Analysis System* (SAS). Hasil kajian menunjukkan bahawa campuran *L. lecanii* + sabun paling efektif menyebabkan kesan mortaliti *P. jackbeardsleyi*, kemudian *L. lecanii* dan *B. bassiana* + sabun, sementara yang terendah adalah *B. bassiana*. Berdasarkan kajian ini, penggunaan fungi entomopatogenik adalah efektif terhadap kutu putih *P. jackbeardsleyi* pada rambutan.

Kata kunci: Patogenitas, *Lecanicillium lecanii*, *Beauveria bassiana*, *Pseudococcus jackbeardsleyi*, rambutan.

INTRODUCTION

Rambutan (*Nephelium lappaceum* L.) is a tropical fruit plant grown in the South East Asian region mainly in Indonesia. The fruit is of a great nutritious value to human health when consumed fresh or as processed fruit. In Indonesia, rambutan is one of the exported fruits and therefore serves a great economic value to the country. The exports of rambutan from Indonesia occupy the second largest position after Thailand (Silitonga 2000). Indonesia's rambutan exports were 47.1 tonnes in 2012 and increased to 66.8 tonnes in 2015 (Indonesia Ministry of Agriculture 2015).

One of the main problems faced by farmers in rambutan cultivation is the insect pests. The presence of insect pests in rambutan affects quality of the fruit and also reduce amount of fruits harvested by farmers. Some important insect pests that attack the rambutan plants including mealybug, *P. jackbeardsleyi* (Hemiptera: Pseudococcidae), *Parasa lepida* (Lepidoptera: Limacodidae), *Adoxophyes privatana* (Lepidoptera: Tortiricidae), *Conopomorpha cramerella* (Lepidoptera: Limacodidae), and *Adoxophyes privatana* (Lepidoptera: Tortiricidae) (Sobir & Martini 2014).

Pseudococcus sp. (Hemiptera: Pseudococcidae) is the commonest amongst all the insect pests that attack rambutan. The pest is widely spread throughout America, Africa, and Asian countries including Taiwan, India, Indonesia, Malaysia, Philippines, Thailand, Singapore, Sri Lanka, and Vietnam (Anura et al. 2012; Ben-Dov et al. 2006; Mani et al. 2013; N'Guessan et al. 2014). The mealybug *P. jackbeardsleyi* is passed to the market commodities by passive dispersal (Williams 2004). It is a big obstacle to the export of fruits because it can survive in postharvest commodities. This is because the mealybug has a layer of white wax like flour all over the body. The mealybug is polyphagous and has several alternative host plants including *Annona* spp. (Magnoliales: Annonaceae), tomato, *Lycopersicon* spp. (Solanales: Solanaceae), banana, *Musa* spp. (Zingiberales: Musaceae), rambutan, *Nephelium lappaceum* L. (Sapindales: Sapindaceae), durian, *Durio zibethinus* Murray (Malvales: Bombacaceae,) pineapple, *Ananas comosus* (L.) (Poales: Bromeliaceae) and rose mallows, *Hibiscus* spp. (Malvales: Malvaceae) (Ben-Dov et al. 2006).

The mealybug is included in the New Zealand government quarantine list of fruits shipped from Indonesia (Biosecurity Act 2014). To overcome the effects of mealybug pest on plant fruits in Indonesia, farmers have always used chemical control measure which involves the use of pesticides such as Imidacloprid. According to Sartiami et al. (2009), Imidacloprid can reduce the population of mealybug on papaya by 40% after four applications, and 60% if the

application of Imidacloprid is combined with soapy water. Other chemical used to control mealybug long time ago was methyl bromide (Hansen et al. 2000). Although several chemicals have been used to control mealybug, little success has been achieved because the mealybugs have a thick waxy layer that is secreted to cover their body, hence resisting the effect from the chemical substances.

Besides the ineffectiveness of chemicals in control mealybug, the application of synthetic pesticides also has a negative impact on the environment and human health, either hence or so we need for an alternative control that is both environmental friendly and cause no harm to human health. One of the alternative measures is the use of entomopathogenic fungi (Ginting et al. 2020). The use of entomopathogenic fungi has several advantages compared to other biological agents, for example they are able to penetrate the body of insects and epizootic in nature. FitzGerald (2014) explains that isolates of *Metarhizium anisopliae* and *B. bassiana* have the potential to be developed as biological control agents to control mealybugs (*Planococcus citri*) and thrips (*Scirtothrips aurantii*) in citrus plants.

The virulence of *B. bassiana* is higher than *M. anisopliae* and *L. lecanii* at the concentration of 5×10^7 conidia/ml, and can cause 98% mortality of *P. ficus* (Signoret) (Hemiptera: Pseudococcidae) (Mohamed 2016). *Beauveria bassiana* caused mortality of *P. marginatus* Williams and Granara de Willink faster than *M. anisopliae* and *V. lecanii*, due to faster penetration and colonization rates. Panyasiri et al. (2007) reported the correlation between the concentration of *M. anisopliae* and the percentage mortality of mealybug, *Pseudococcus cryptus* (Hempel). However, the information on pathogenicity of entomopathogenic fungi *L. lecanii* and *B. bassiana* against mealybug in rambutan has not been reported. Based on this description, this study aims to examine the pathogenicity of entomopathogenic fungi *L. lecanii* and *B. bassiana* against mealybug *P. jackbeardsleyi* in rambutan.

METRIALS AND METHODS

Study Site

This study was conducted from February until November 2019. The mealybugs, *P. jackbeardsleyi* were sampled from rambutan trees at Timur Indah, Bengkulu City, Indonesia and brought back to the Plant Protection Laboratory, Department of Plant Protection, Faculty of Agriculture, Universitas Bengkulu for identification process and bioassay.

Species Identification

The *P. jackbeardsleyi* species was identified using species key (Williams & Granara de Willink 1992) before they were analysed for bioassay using entomopathogenic fungi.

Identification of Entomopathogenic Fungi

The entomopathogenic fungus used was a collection of the Plant Protection Laboratory, Department of Plant Protection, Faculty of Agriculture, Bengkulu University. Morphological identification of the fungi was carried out using Humber (1997). The optical microscope was used to make observations on the morphological features of the fungi colonies grown on Potato Dextrose Agar (PDA) media in petri dishes. The colony features observed include colony's color, shape, texture and edge shape. Other observations made include the forms of conidia and the presence of hyphae.

Preparation of the *B. bassiana* and *L. lecanii*

Fungi isolates of *B. bassiana* and *L. lecanii* were used for insect bioassay. The isolates were prepared from the entomopathogenic fungi grown on petri dishes with Potato Dextrose Agar (PDA) (potato 200g, dextrose 20g, agar 15g, chloramphenicol 0.5g, and aquadest 1L) at 21°C. After 3 weeks, the fungi conidia were harvested, by adding 10 ml of sterilized aquades to the fungi colonies to form a suspension. Then 0.1% Triton X-100 was added to the suspension as an adhesive and grading agent. Its density was calculated using hemocytometer.

Pathogenicity of *B. bassiana* and *L. lecanii* against Nymph of *P. jackbeardsleyi*

Second instar of *P. jackbeardsleyi* nymphs were used for bioassay. A total of 5 nymphs of *P. jackbeardsleyi* were placed in a plastic container with a diameter of 6.5 cm and height of 4.5 cm and fed with rambutan fruit. Concentration of entomopathogenic fungi use for bioassay is shown in Table 1 (*B. bassiana* (Bb): Bb 10⁶, Bb 10⁷, Bb 10⁸ (conidia/ml), Bb 10⁶ + Soap 0.2%, Bb 10⁷ + Soap 0.2%, Bb 10⁸ + Soap 0.2%; *Lecanicillium lecanii* (LL): LL10⁶, LL10⁷, LL10⁸ (conidia/ml), LL 10⁶ + Soap 0.2%, LL 10⁷ + Soap 0.2%, LL 10⁸ + Soap 0.2%, soap 0.2%, and control (sterile water)). The soap used in the test was Benzalkonium Chloride. Each treatment was repeated three times. Fungal application was carried out by spraying nymphs with a conidia suspension of 10 ml (Tefera & Pringle 2003). Larval mortalities were recorded every day for five days after application.

Data Analysis

Data obtained were analyzed using the SAS program version 6.2. Duncan's Multiple Range Test at the 5% significance level was performed if there were differences between treatments. Determination of lethal concentration (LC); LC₅₀, LC₇₅, and lethal time (LT); LT₅₀, LT₇₅ values was carried out by Probit Analysis (Finney 1971).

RESULTS

Mortality of *P. jackbeardsleyi* Nymphs after Treatments

All treatments caused mortality of *P. jackbeardsleyi* nymphs under laboratory conditions. The highest cumulative mortality rate of *P. jackbeardsleyi* nymphs recorded was 93% and 100% at concentrations of 10⁷ and 10⁸ (conidia/ml), respectively for entomopathogenic fungi treatments; *B. bassiana*, *B. bassiana* + soap, and *L. lecanii* + soap, and *L. lecanii* (Table 1).

Table 1 Cumulative mortality of *P. jackbeardsleyi* for five days after treatment with various concentrations of entomopathogenic fungi *B. basianna* and *L. lecanii*, and fungi mixture with soap.

Treatment	% Mortality (±SD)
Control (sterile water)	0±0.00a
Soap	13±0.67a
Bb 10 ⁶ (conidia/ml)	53±0.58b
Bb 10 ⁷ (conidia/ml)	73±1.15b
Bb 10 ⁸ (conidia/ml)	93±0.58d
Bb 10 ⁶ (conidia/ml) + Soap	80±1.00c
Bb 10 ⁷ (conidia/ml) + Soap	87±0.58c
Bb 10 ⁸ (conidia/ml) + Soap	100±0.00e
LL 10 ⁶ (conidia/ml)	67±0.58b
LL 10 ⁷ (conidia/ml)	73±0.58b

LL 10 ⁸ (conidia/ml)	93±0.58d
LL 10 ⁶ (conidia/ml) + Soap	80±1.00c
LL 10 ⁷ (conidia/ml) + Soap	93±0.58d
LL10 ⁸ (conidia/ml) + Soap	100±0.00e

*The number followed by the same letter in the same column is not significantly different according to Duncan's test at 5% significance level. Bb: *B. basianna*, LL: *L. lecanii*.

*Soap: Benzalkonium chloride 0.2%




Figure 1 The cumulative mortality of *P. jackbeardsleyi* nymphs against treatment of various with various concentrations of entomopathogenic fungi, *B. basianna* and *L. lecanii*, and fungi mixture with soap

Lethal Concentration LC₅₀ and LC₇₅ of Treatments

The density of the conidia of fungus was used to determine the mortality of *P. jackbeardsleyi* nymphs. Based on the results of Probit Analysis on the fifth day observation, treatment with the LC₅₀ and LC₇₅ was found *L. lecanii* + soap, followed by *L. lecanii*, *B. bassiana* + soap, *B. bassiana* (Table 2).

Table 2. LC₅₀ and LC₇₅ values of each treatment of entomopathogenic fungi, *B. basianna*, *L. lecanii*, and fungi mixture with soap

Treatments	LC ₅₀ (conidia/ml)	LC ₇₅ (conidia/ml)
<i>Beauveria bassiana</i>	21.6 x 10 ⁸	2.83 x 10 ⁹
<i>Beauveria bassiana</i> + soap	20.0 x 10 ⁸	2.74 x 10 ⁹
<i>Lecanicillium lecanii</i>	17.2 x 10 ⁸	1.91 x 10 ⁹
<i>Lecanicillium lecanii</i> + soap	16.8 x 10 ⁸	1.87 x 10 ⁹

Lethal Time LT_{50} and LT_{75} of Treatments

The ability of the treatments to cause mortality of *P. jackbeardsleyi* nymphs was shown by LT_{50} and LT_{75} values for each treatment against *P. jackbeardsleyi* nymphs at a density 10^8 (conidia/ml). The results showed that mixture of *B. bassiana* and *L. lecanii* with soap as treatment gave the lowest lethal time values, LT_{50} (3.5 days) and LT_{75} (3.9 days). These treatments were followed by *L. lecanii*, and *B. bassiana* (Table 3).

Table 3 LC_{50} , LC_{75} values of with various concentrations of entomopathogenic fungi, *B. bassiana* and *L. lecanii*, and fungi mixture with soap against *P. jackbeardsleyi* nymphs

Treatments	LT_{50} (Day)	LT_{75} (Day)
<i>Beauveria bassiana</i>	3.8	4.2
<i>Beauveria bassiana</i> + soap	3.5	3.9
<i>Lecanicillium lecanii</i>	3.6	4.05
<i>Lecanicillium lecanii</i> + soap	3.5	3.9

Growth of white mycelia after mortality of *P. jackbeardsleyi* nymphs

Observation through the microscope shown growth of white mycelia from *P. jackbeardsleyi* nymphs' cadavers (Figure 2).




Figure 2. a) White mycelia formed on *P. jackbeardsleyi* nymphs cadavers by a, *B. bassiana*; b), *L. lecanii*.

DISCUSSION

The difference in concentration of treatments caused the differences in conidia density hence differences in percentage mortality of *P. jackbeardsleyi* nymph. The treatments with high conidia density gave greater opportunities for conidia to penetrate into the body of *P. jackbeardsleyi* nymphs and cause mortality. The highest *P. jackbeardsleyi* nymph mortality was at the conidia density $Bb 10^8 + 0.2\%$ soap and $LL 10^8 + 0.2\%$ soap which reach 100% mortality. The lowest *P. jackbeardsleyi* nymph mortality was with $Bb 10^6$ treatment. These results correspond with the findings from a number of studies. Shinde et al. (2010) explains that one of the factors of entomopathogenic fungal infections in insects is the number of inoculums. Inglis et al. (2001) reported the relationship between the virulence of *B. bassiana*, *M. anisopliae*, and the

concentration of entomopathogenic fungi for insect pests. The higher conidia density was applied, so the chance of conidia contacts greater with pests, that it gives more opportunities to penetrate the pest body. Chavan and Kadam (2009) also reported that mortality of mealybug *M. hirsutus* nymphs was 82.5% with *L. lecanii* application. Furthermore, Jeyarani et al. (2011) reported that *B. bassiana* was effective than *Cladosporium cladosporioides* (Fresenius) against papaya mealybug, *Paracoccus marginatus*. Moreover, *B. bassiana* isolated from Bengkulu caused 76.7%-80.6% mortality on *Hyphotenemus hampei* Ferrari when applied at 10^9 conidia/ml (Apriyanto & Nadrawati 2019). Agustin (2014) reported that *L. lecanii* at a density of 10^8 conidia/ml, was able to cause 67.62% mortality of *O. furnacalis* larvae, while at a density of 10^9 conidia/ml the mortality rate was 71.25%.

The lowest lethal concentration ($LC_{50,75}$) was found in the treatment of *L. lecanii* + soap. This was followed by the treatment of *L. lecanii*, *B. bassiana* + soap, *B. bassiana*, and soap. These results showed that the treatment of *L. lecanii* + soap was the most effective against mealy bug. These results correspond with the findings by Bhadani et al. (2017), which showed that *L. lecanii* 2.0 gm/l + profenophos 50 EC 0.025% and *L. lecanii* @ 2.0 gm/l + flonicamid 50 WG 0.0125% was more effective to reduce the population of mealy bugs, *Maconellicoccus hirsutus* in apples both under laboratory and field conditions with mortality of 68.14-72.69%, on the fifth day, than *B. bassiana* 2.0 gm/l + profenophos 50 EC 0.025% @ 0.5ml/l and *B. bassiana* 2.0 gm/l + flonicamid 50 WG 0.0125% @ 0.25 gm/l with mortality of 44.40-51.23%. Leland et al. (2005), explains that, the differences in pathogenicity of different entomopathogenic fungi strains is influenced by the physiological and enzymatic properties of each isolate. The pathogenesis of entomopathogenic fungi depends on the pathogen and host, whereby the cuticle of host affects the infection process, adhesion, germination and differentiation of aspersorium (Vega & Kaya 2012). The lethal concentration also range of a fungus isolate depends on the strain, the host insect, and the application method (Baidoo & Ackuaku 2011).

Entomopathogenic fungi function in several ways during control of insect populations. During the process of infection, the entomopathogenic fungi produce a number of enzymes, such as proteases, chitinases, and lipases that function to degrade the host cuticle and facilitate attachment of the entomopathogenic fungus conidia to the cuticle of the host. The entomopathogenic fungi also produce several types of toxins (e.g. *B. bassiana* produces bassianin, beauvericin, bassianolide, beaverolides and tenellin while *L. lecanii* produces dipcolonic acid, hydroxy carboxylic acid and cyclosporine) which cause an increase in the pH of haemolymph, clotting and stopping the circulation of haemolymph in insect. As a result, the function of the insect's hemolymph and nucleus are interfered, resulting in death of the infected insect due to swelling and hardening (Vey et al. 2001).

The results of the lethal time values showing the time taken by the entomopathogenic fungi (*B. bassiana* and *L. lecanii*) treatments to cause mortality of the *P. jackbeardsleyi* nymphs showed that *B. bassiana* + soap and *L. lecanii* + soap treatments had the lowest LT_{50} value of 3.5 days each, followed by *L. lecanii* (3.6 days) and then *B. bassiana* (3.8 days). The LT_{50} value of *B. bassiana* treatment different with findings by Ahmed (2013), which showed that the application of *B. bassiana* strain PDRL1187 against *Phenacoccus solenopsis* has LT_{50} value of 13.4 days, and *B. bassiana* PDRL1147 has LT_{50} 13.9 days.

The lethal time values obtained were related to isolate virulence and host susceptibility in a way that the isolate with higher virulence took less time than the isolate with lower virulence to kill the host (Vega & Kaya 2012). Inglis et al. (2001) explains that the difference in virulence of the isolate is affected by the process of fungus infecting insects starting from attaching of

conidia, germination, penetration, invasion and colonization in haemocytes, tissues, and organs of the host. (Vega & Kaya 2012) explains that the effectiveness of entomopathogenic fungi begins from 48 hours post inoculation when the hyphae penetrates the integument in the trachea, epithelium, and epidermal cells. At the 72 hours, the fat tissue is damaged and after 96 hours the mortality of the host occurs.

CONCLUSION

The entomopathogenic fungi treatment of *L. lecanii* + soap is the most effective, followed by *L. lecanii*, *B. bassiana* + soap and *B. bassiana* against *P. jackbeardsleyi*. This is because *L. lecanii* + soap had the lowest LC₅₀ and LC₇₅ value amongst all the treatments tested. Also, *L. lecanii* + soap and *B. bassiana* + soap treatments had the lowest lethal time value compared to others. Therefore, it can be concluded that entomopathogenic fungi *L. lecanii* and *B. bassiana* tested are appropriate control alternatives against *P. jackbeardsleyi* in rambutan.

REFERENCES

- Ahmed, A. 2013. Use of entomopathogenic fungi in biological control of cotton mealybug (*Phenacoccus solenopsis*) and mustard aphid (*Lipaphis erysimi*). Thesis Master. Department of Botany, University of Karachi, Pakistan.
- Agustin, D. 2014. The effectiveness of the entomopathogenic fungi *Beauveria bassiana* (Balsamo) Vuillemin and *Lacanicillium lecanii* (Zimm.) Zare & Gams against asian corn stem borer *Ostrinia furnacalis* Guenee (Lepidoptera: Crambidae). Thesis Master. Bogor Agricultural University, Bogor.
- Anura, U.G., Sirisena, I., Watson, G.W., Hemachandra, K.S. & Wijayagunasekara, H.N.P. 2012. Mealybugs (Hemiptera: Pseudococcidae) introduced recently to Sri Lanka, with three new country records. *Pan-Pacific Entomologist* 88(3): 365-367.
- Apriyanto, D., & Nadrawati. 2019. Laboratory evaluation of Bengkulu isolates of *Beauveria bassiana* and *Metarhizium anisopliae* against coffee berry borer, *Hyphotenemus hampei*, using spraying method. *J. HPT Tropika* 19(2): 93-100.
- Baidoo, P.K. & Ackuaku, S.K. 2011. The effects of spore concentration of entomopathogenic fungi on larval mortality and development of the maize stem borer, *Eldana saccharina* (Lepidoptera: Pyralidae). *Journal of Applied Biosciences* 47: 3221-3229.
- Ben-Dov Y., Miller, D.R. & Gibson, G.A.P. 2006. ScaleNet, Life Histories. <http://www.sel.barc.usda.gov/scalenet/lifehist.htm>. [20 August 2019].
- Bhadani, D.J., Kabaria, B.B. & Ghelani, M.K. 2017. Bio-efficacy of entomopathogenic fungi against mealybug, *Maconellicoccus hirsutus* (Green) infesting custard apple in Junagadh. *Journal of Entomology and Zoology Studies* 5(5): 285-289.
- Biosecurity Act. 2014. Import health standard fresh fruit/vegetables Mangosteen (*Garcinia mangostana*) from Indonesia. http://52.64.242.214/uploads/PIAT_content/pdfs/IHS/Mangosteen-%20Indo_IHS.pdf. [15 August 2019].
- Chavan, B.P. & Kadam, J.R. 2009. Effect of combination of adjuvants on liquid formulations of *Verticillium lecanii* (Zimmermann) Viegas and their efficacy. *Journal Biological Control* 23: 73-77.
- Finney, D.J. 1971. *Probit Analysis*. 3rd Edition. New York (US): Cambridge University Press.
- FitzGerald, V.C. 2014. Screening of entomopathogenic fungi against citrus mealybug (*Planococcus citri* (Risso)) and citrus thrips (*Scirtothrips aurantii* Faure). Thesis Master. Rhodes University, South Africa.
- Ginting, S., Zarkani, A., Wibowo, R.H. & Sipriyadi. 2020. New invasive pest *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) attack on varieties of corn in Bengkulu. *Serangga* 25(1): 104-115.

- Hansen, J.D., Drake, S.R., Moffitt, H.R., Albano, D.J. & Heidt, M.L. 2000. Methylbromide fumigation of five cultivars of sweet cherries as a quarantine treatment against codlingmoth. *HortTechnology* 10(1): 194-198.
- Humber, R.A. 1997. *Manual of Techniques in Insect Pathology: Identification*. California (US): Academic Press.
- Indonesia Ministry of Agriculture. 2015. Rambutan fruit exports in 2012-2015. <https://aplikasi.pertanian.go.id/bdsp/hasil-ind.asp> [15 April 2019].
- Inglis, G.D, Goettel, M.S., Butt, T.M. & Strasser. H. 2001. Use of hyphomycetous fungi for managing insect pests. In Butt, T.M., Jackson, C.W. & Magan, N. (eds.). *Fungi as Biocontrol Agents: Progress, Problems and Potential*, pp. 23–69. Wallingford, United Kingdom: CAB International/AAFC.
- Jeyarani, S., Banu, J.G. & Ramaraju, K. 2011. First record of natural occurrence of *Cladosporium cladosporioides* (Fresenius) de Vries and *Beauveria bassiana* (Bals.-Criv.) vuill on papaya mealybug, *Paracoccus marginatus* Williams from India. *Journal of Entomology* 8: 274-279.
- Leland, E.J., Mcguire, M.R., Grace, G.A., Jaronski, S.T., Ulloa M., Park, Y.H. & Plattner, R.D. 2005. Strain selection of fungal entomopathogen, *Beauveria bassiana*, for control of plant bugs (*Lygus* spp.) (Heteroptera: Miridae). *Biological Control* 35: 104–114.
- Mani, M., Joshi, S., Kalyanasundaram, M., Shivaraju, C., Krishnamoorthy, A., Asokan, R. & Rebijith, K.B. 2013. A new invasive Jack Beardsley Mealybug, *Pseudococcus jackbeardsleyi* (Hemiptera: Pseudococcidae) on papaya in India. *Florida Entomologist* 96(1): 242-245.
- Mohamed, G.S. 2016. Virulence of entomopathogenic fungi against the vine mealybug, *Planococcus ficus* (Signoret) (Hemiptera: Pseudococcidae). *Egyptian Journal of Biological Pest Control* 26(1): 47-51
- N'Guessan, P.W., Watson, G.W., Brown, J.K. & N'Guessan, F.K. 2014. First record of *Pseudococcus jackbeardsleyi* (Hemiptera: Pseudococcidae) from Africa, Côte D'Ivoire. *Florida Entomologist* 97(4): 1690-1693.
- Panyasiri, C., Attathom, T. & Poehling, H.M. 2007. Pathogenicity of entomopathogenic fungi-potential candidates to control insect pests on tomato under protected cultivation in Thailand. *Journal of Plant Diseases and Protection* 114 (6): 278–287.
- Sartiarni, D, Pudjianto, & Buchori, D. 2009. Strengthening local natural enemies of papaya whitefly infestation (*Paracoccus marginatus*). Faculty of Agriculture, Bogor Agricultural University, Bogor. Final report.
- Shinde, S.V, Patel, K.G, Purohit, M.S, Pandya, J.R & Sabalpara. 2010. *Lecanicillium lecanii* (Zimm.) Zare and Games an important biocontrol agents for the management of insect pests-A review. *Agr Review* 31(4): 235-252.

- Silitonga, D. 2000. Analysis of the financial feasibility of rambutan farming investment in the Binjai Barat sub-district. Thesis Master. Bogor Agricultural University, Bogor.
- Sobir & Martini, E. 2014. *Guidelines for Cultivating Durian and Rambutan in Mixed Gardens*. Bogor: Southeast Asia Regional Program World Agroforestry Center.
- Tefera, T. & Pringle, K.L. 2003. Effect of exposure method to *Beauveria bassiana* and conidia concentration on mortality, mycosis, and sporulation in cadavers of *Chilo partellus* (Lepidoptera: Pyralidae). *Journal of Invertebrate Pathology* 84(2):90–95.
- Vega, F. & Kaya, H. 2012. *Insect Pathology*. New York (US): Academic Press.
- Vey, A., Hoagland, R.E. & Butt, T.M. 2001. Toxic metabolites of fungal biocontrol agents. In Butt, T.M, Jackson, C. & Magan, N. (eds.). *Fungi as biocontrol agents. Progress, problems and potential*, pp. 311-346. Oxford (US): CAB International.
- Williams, D.J. & Granara de Willink, M.C. 1992. *Mealybugs of Central and South America*. Wallingford (US): CAB International.
- Williams, D.J. 2004. *Mealybugs of Southern Asia*. Kuala Lumpur: United Selangor Press.